

## Inhibition of the helicase activity of HCV NTPase/helicase by 1- $\beta$ -D-ribofuranosyl-1,2,4-triazole-3-carboxamide-5'-triphosphate (ribavirin-TP)

Peter Borowski<sup>1</sup>, Melanie Lang<sup>1</sup>, Andreas Niebuhr<sup>1</sup>, Annemarie Haag<sup>1</sup>,  
Herbert Schmitz<sup>1</sup>, Julian Schulze zur Wiesch<sup>2</sup>, Joonho Choe<sup>3</sup>,  
Maria A. Siwecka<sup>4</sup> and Tadeusz Kulikowski<sup>4</sup>✉

<sup>1</sup>*Abteilung für Virologie, Bernhard-Nocht-Institut für Tropenmedizin, Hamburg, Germany;*  
<sup>2</sup>*Klinik und Poliklinik Innere Medizin, Universitätsklinikum Hamburg-Eppendorf, Hamburg, Germany;*  
<sup>3</sup>*Department of Biological Sciences, Korea Advanced Institute of Science and Technology, Taejeon, Korea;*  
<sup>4</sup>*Institute of Biochemistry and Biophysics, Polish Academy of Sciences, Warszawa, Poland*

Received: 8 January, 2001; revised: 4 April, 2001; accepted: 4 July, 2001

**Key words:** HCV helicase, enzyme inhibition, ribavirin-5'-triphosphate

**In the presented study the ribavirin-TP – an established inhibitor of the NTPase activity of the superfamily NTPase/helicases II – was investigated as an inhibitor of the unwinding activity of the hepatitis C virus (HCV) NTPase/helicase. The kinetics of the reaction revealed that ribavirin-TP reduces the turnover number of the helicase reaction by a mechanism that does not correspond to that of the inhibition of the NTPase activity. Our results suggest that derivatives of ribavirin-TP with enhanced stability towards hydrolytic attack may be effective inhibitors of the enzyme.**

NTPase/helicases unwind duplex RNA or DNA structures by disrupting the hydrogen bonds that keep the two strands together [1–3]. This reaction is coupled with the hydrolysis of  $\gamma$ -phosphate of ATP. All NTPase/heli-

cases known to date display ATPase activity [2, 3]. Recently, it was found that, in the presence of a nonhydrolyzable ATP analogue, adenosine-5'-(3-thiotriphosphate) (ATP- $\gamma$ -S), only a low level of unwinding of HCV dsRNA

---

✉To whom correspondence should be addressed: Institute of Biochemistry and Biophysics, Polish Academy of Sciences, A. Pawińskiego 5a, 02-106 Warszawa, Poland; e-mail: tk@ibbrain.ibb.waw.pl

**Abbreviations:** ATP- $\gamma$ -S, adenosine-5'-(3-thiotriphosphate); HCV, hepatitis C virus; NDP, nucleoside-5'-diphosphate(s); NMP, nucleoside-5'-monophosphate(s); NS, nonstructural proteins; NTP, nucleoside-5'-triphosphate(s); NTPase, nucleoside triphosphatase, ribavirin-TP, 1- $\beta$ -D-ribofuranosyl-1,2,4-triazole-3-carboxamide-5'-triphosphate; TBE, Tris/borate/EDTA buffer.

could be detected [4]. The NS3 – like proteins of flaviviruses, pestiviruses and of HCV belong to superfamily II (SFII) [2]. The ATP-binding site of the NTPase/helicase contains conserved amino-acid sequences called Walker motifs A and B that are located on the surface of domain I of the three-domain enzymes [5]. These motifs are responsible for the binding of the terminal phosphate groups of ATP and the  $Mg^{2+}$  of the Mg-ATP complex respectively [5–7]. Mutations in these motifs eliminate the ATP hydrolytic activity of the enzymes [6, 8].

On the basis of these results one could speculate that a reduction of the accessibility of the ATP-binding site for ATP may lead to inhibition of the ATPase and, consequently, of the helicase activity of the NTPase/helicases. Indeed, in our previous studies we have demonstrated that compounds that reduced the binding of ATP to proteolytically isolated domain I of the HCV NTPase/helicase are capable of inhibiting the ATPase activity of the enzyme [9]. Nevertheless, there is no simple correlation between the turnover number of the NTPase reaction and the number of helicase unwinding cycles of the enzymes. Further analysis using a West Nile (WN) virus NTPase/helicase, a further SFII enzyme, revealed that O<sup>6</sup>-benzylguanine and N-chloroethylguanine derivatives modulated, i.e. enhanced or reduced, the ATPase activity of the enzyme without influencing the helicase activity. Moreover, the helicase activity may be up- or down-regulated without corresponding changes of the requirement for ATP [10].

Therefore we performed a study, presented here, in which the inhibitory potential of 1- $\beta$ -D-ribofuranosyl-1,2,4-triazole-3-carboxamide-5'-triphosphate (ribavirin-TP) towards the helicase activity of HCV NTPase/helicase was tested. This compound was established to be a competitive (with regard to ATP) inhibitor of the NTPase activity of HCV and WN virus NTPase/helicases [10, 11].

## MATERIALS AND METHODS

***Production and purification of the HCV NTPase/helicase.*** HCV NTPase/helicase was expressed in *Escherichia coli* and purified as previously described [12, 13]. The purity of the obtained enzyme was around 85%. The purification procedure was completed by gel exclusion chromatography on Superdex 200 (Amersham-Pharmacia) [11]. The final preparation contained homogeneous HCV NTPase/helicase.

***ATPase and helicase assays.*** A standard ATPase assay was performed according to the charcoal adsorption method described previously [9, 11]. The helicase activity of the enzyme was determined using a DNA substrate that was obtained by annealing two partly complementary DNA oligonucleotides. The synthesised oligonucleotides had sequences corresponding to the deoxynucleotide version of the RNA strands described previously [4, 10].

***Synthesis of 1- $\beta$ -D-ribofuranosyl-1,2,4-triazole-3-carboxamide-5'-triphosphate (ribavirin-TP).*** Ribavirin-TP was synthesised according to the modified Yoshikawa-Ludwig-Mishra-Broom procedure [11, 14–17].

***Other methods.*** Kinetic parameters of the reactions were determined by non-linear-regression analysis using ENZFITTER (BioSoft) and SIGMA PLOT (Jandel Corp.). Protein concentration was measured by the method of Lowry *et al.* [18]. The purity of the proteins was determined by densitometric analysis of the SDS/polyacrylamide gels stained with Coomassie Blue [19].

## RESULTS AND DISCUSSION

### Purity of the enzyme

It is known that the NTPase and helicase activities of different preparations of HCV en-

zymes expressed in *E. coli* differ in their specific activity, the response to polynucleotides or their affinity to substrate [11]. Since the discrepancies between the data result often from differences in the purity of the final preparations or various reaction conditions, the HCV NTPase/helicase used in this study was purified so as to obtain a homogeneous enzyme preparation and optimised conditions of the reaction had to be established.


### Substrates and kinetic parameters of HCV NTPase/helicase

The NTPase activity of the enzyme was assayed by a previously described charcoal adsorption method [9, 11]. Direct comparison of the kinetic parameters obtained under optimised conditions showed no significant differences between the enzyme preparation used here and that used in previous studies [9, 11]. The two enzymes displayed similar temperature optima, time course of the reaction and dependency on divalent ions. However, the  $K_m$  value for ATP estimated for the enzyme used here was  $143.5 \mu\text{M}$ , in contrast to  $11.0 \mu\text{M}$  for the enzyme described by us previously [9, 11].

The helicase activity of the enzyme was investigated with a partly double-stranded DNA substrate well characterised in previous studies. The substrate was applied at non-saturating concentrations [4, 10]. When the kinetic parameters of the unwinding and ATPase activities of the enzyme were compared, some discrepancies between the optimum conditions were observed. The most prominent difference, important for the further study, probably due to the strongly limited concentrations of the DNA-substrate, is that the half-optimum ATP concentration for the helicase turnover was 5 fM (Borowski, P., Niebuhr, A. & Schmitz, H., unpublished data).

An important feature of the unwinding reaction is that at the ATP concentration corresponding to the  $K_m$  value for the ATPase reaction ( $143.5 \mu\text{M}$ ) and with 4.7 pM of the

DNA-substrate (nucleotide base), the plateau of the unwinding reaction was reached after 20 min (Fig. 1). There were no significant differences when the helicase and ATPase activities were compared in respect of their dependency on divalent ions and temperature optimum (not shown).



**Figure 1. Time course of the unwinding reaction mediated by HCV NTPase/helicase.**

The helicase assays were performed with 10 pmol of the enzyme and the unwinding reaction was terminated at the times indicated in the figure as described under Materials and Methods. The samples were separated on a 15% polyacrylamide TBE gel containing 0.1% SDS and the dried gel was exposed to X-ray film at  $-80^{\circ}\text{C}$  for 14 h. The autoradiography shown is representative of three independent experiments.

### Response of the HCV NTPase/helicase to ribavirin and ribavirin-TP


Previous studies indicated that, at a very low ATP concentration, i.e.  $<1/1000$  of  $K_m$  values, ribavirin added at a concentration range of  $50-750 \mu\text{M}$  could act as a competitive ATPase inhibitor [11]. Ribavirin *versus* ATP titration curves did not reveal, however, any inhibitory potential of the compound towards the helicase activity of the enzyme. Similar results were obtained previously with the WN virus NTPase/helicase [10] and with the Japan encephalitis virus (JEV) enzyme, belonging to the SFII NTPase/helicases too (not shown).

It was found previously that ribavirin-TP exhibited a higher inhibitory effect on the

NTPase activity of the HCV and WN virus enzymes than did ribavirin [11]. Thus, one could speculate that an effective inhibition of the NTPase activity can reduce the energy supply and therefore inhibit the unwinding activity of the enzyme. The helicase activity of the enzyme was investigated as a function of increasing concentrations of ribavirin-TP and the reaction was terminated after 20 min, i.e. at a time at which >95% of the DNA-substrate was unwound. As expected, when the ATP concentration was adjusted to a concentration equal to the  $K_m$  value, only a moderate inhibi-

extent. Thus, a reduction of the reaction time to 10 min at which about 60% of DNA-substrate was unwound resulted in  $IC_{50}$  of  $90 \mu M$  ( $\pm 10$ ). When the reaction time was limited to 5 min (35% of substrate unwound) the  $IC_{50}$  was  $30 \mu M$  ( $\pm 13$ ). Nevertheless, the inhibition was not complete and reached 35% of the control at ribavirin-TP concentrations corresponding to a values 3–5 times as high as the  $IC_{50}$  i.e.  $200 \mu M$  ( $\pm 8.5$ ) (Fig. 2).

These data are obviously contrary to the previous results obtained with the WN virus NTPase/helicase [10]. In the case of the WN



**Figure 2.** Inhibition of the unwinding reaction mediated by HCV NTPase/helicase and its dependency on the duration of the reaction.

The helicase assays were performed as described above in the presence of increasing concentrations of ribavirin-TP. The reaction was terminated after 5 min (▼), 10 min (●) and 20 min (■) duration of the reaction. The samples were proceeded as described in the legend to Fig. 1 and the helicase activity was quantified according to the procedure reported previously [10]. The obtained values are presented as percentages of controls (no inhibitor added) and are averages of three independent assays.

tion with an  $IC_{50}$  of  $150$ – $200 \mu M$  was seen. The inhibition was not complete and reached a maximum of 35% of the control at ribavirin-TP concentrations higher than  $300 \mu M$  ( $\pm 10$ ). Surprisingly, a similar extent of inhibition was also observed when ATP concentration was reduced to  $0.01 \mu M$  or  $5 \text{ fM}$ , i.e. to concentrations at which ribavirin-TP acts as a potent inhibitor of the ATPase activity of the enzyme (ref. [11] and unpublished data). Interestingly, the inhibitory potential of the compound was higher when the reaction was terminated earlier, i.e. at times at which the DNA-substrate was unwound only to some

virus enzyme no apparent correlation between the duration of the reaction and inhibitory potential of the ribavirin-TP was seen (see ref. [10] and unpublished data). One possible explanation for this kinetic behaviour is the relatively low specificity of the HCV enzyme for NTP [20, 21]. Thus it could not be ruled out that the enzyme hydrolysed the ribavirin-TP to less potent derivatives. Indeed, our preliminary data indicate that the HCV enzyme produces not only NDP, but also substantial amounts of NMP in the course of hydrolysis of ATP and other NTPs. Moreover, the products of the hydrolysis were not capa-

ble inhibiting the helicase activity of the enzyme (Borowski, P., Niebuhr, A., Schmitz, H., unpublished data)

In this context, it is apparent that introduction of  $\beta$ - and  $\gamma$ -phosphate groups into the 5'-position of ribavirin molecule could enhance the inhibitory potential of the drug. On the other hand, in the light of the low specificity of the enzyme for phosphate groups of NTPs only the compounds with enhanced resistance to hydrolysis of the terminal phosphate group(s) seem to be suitable as inhibitors of the helicase activity. Investigations on nonhydrolysable analogues of ribavirin-TP are currently under way.

## REFERENCES

- Gorbalenya, A.E. & Koonin, E.V. (1993) Helicases: Amino acid sequence comparisons and structure-function relationships. *Curr. Opin. Struct. Biol.* **3**, 419-429.
- Kadare, G. & Haenni, A. (1997) Virus-encoded RNA helicase. *J. Virol.* **71**, 2583-2590.
- Hodgman, T.C. (1988) A new superfamily of replicative proteins. *Nature* **333**, 22-23.
- Gallinari, P., Brennan, D., Nardi, C., Brunetti, M., Tomei, L., Steinkühler, C. & De Francesco, R. (1998) Multiple enzymatic activities associated with recombinant NS3 of hepatitis C virus. *J. Virol.* **72**, 6758-6769.
- Walker, J.E., Saraste, M., Runswick, M.J. & Gay, N.J. (1982) Distantly related sequences in the alpha- and beta-subunits of ATP synthase, myosin, kinases and other ATP-requiring enzymes and a common nucleotide binding fold. *EMBO J.* **1**, 945-951.
- Kim, J., Morgenstern, K., Griffith, J., Dwyer, M., Thomson, J., Murcko, M., Lin, C. & Caron, P. (1998) Hepatitis C virus NS3 RNA helicase domain with a bound oligonucleotide: The crystal structure provides insights into the mode of unwinding. *Structure* **6**, 89-100.
- Yao, N., Hesson, T., Cable, M., Hong, Z., Kwong, A., Le, H. & Weber, P. (1997) Structure of the hepatitis C virus RNA helicase domain. *Nat. Struct. Biol.* **4**, 463-467.
- Kim, D.W., Kim, J., Gwack, Y., Han, J.H. & Choe, J. (1997) Mutational analysis of the hepatitis C virus RNA helicase. *J. Virol.* **71**, 9400-9404.
- Borowski, P., Kuehl, R., Mueller, O., Hwang, L.-H., Schulze zur Wiesch, J. & Schmitz, H. (1999) Biochemical properties of a minimal functional domain with ATP-binding activity of the NTPase/helicase of hepatitis C virus. *Eur. J. Biochem.* **266**, 715-723.
- Borowski, P., Niebuhr, A., Mueller, O., Bretner, M., Felczak, K., Kulikowski, T. & Schmitz, H. (2001) Purification and characterization of West Nile virus NTPase/helicase. Evidence for dissociation of the NTPase and helicase activities of the enzyme. *J. Virol.* **75**, 3220-3229.
- Borowski, P., Mueller, O., Niebuhr, A., Kalitzky, M., Hwang, L.-H., Schmitz, H., Siwecka, A.M. & Kulikowski, T. (2000) ATP-binding domain of NTPase/helicase as a target for hepatitis C antiviral therapy. *Acta Biochim. Polon.* **47**, 173-180.
- Kim, D.W., Gwack, Y., Han, H.J. & Choe, J. (1995) C-terminal domain of the hepatitis C virus NS3 protein contains an RNA helicase activity. *Biochem. Biophys. Res. Commun.* **215**, 160-166.
- Gwack, Y., Kim, D.W., Han, J. & Choe, J. (1996) Characterization of RNA binding activity and RNA helicase activity of hepatitis C virus NS3 protein. *Biochem. Biophys. Res. Commun.* **225**, 654-659.
- Ludwig, J. (1981) Chemical synthesis of nucleoside triphosphates. *Acta Biochim. Biophys. Acad. Sci. Hung.* **16**, 131-133.
- Bowles, W.A., Schneider, F.H., Lewis, L. & Robins, R.K. (1963) Synthesis and antitumor activity of 9-(tetrahydro-2-furyl)purine analogs

- of biologically important deoxynucleosides. *J. Med. Chem.* **6**, 471-480
- 16.** Mishra, N.C.U. & Broom, A.D. (1991) A novel synthesis of nucleoside 5'-triphosphates. *J. Chem. Soc., Chem. Commun.* 1276-1277.
- 17.** Yoshikawa, M., Kato, T. & Takenishi, T. (1967) A novel method of phosphorylation of nucleosides to 5'-nucleotides. *Tetrahedron Lett.* **50**, 5065-5068.
- 18.** Lowry, O.H., Rosebrough, N.J., Farr, A.J. & Randall, R.J. (1951) Protein measurement with the Folin phenol reagent. *J. Biol. Chem.* **193**, 265-275.
- 19.** Laemmli, U.K. (1970) Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature* **227**, 680-685.
- 20.** Preugschat, F., Averett, D., Clarke, B. & Porter, D. (1996) A steady state and pre-steady-state kinetic analysis of the NTPase activity associated with the hepatitis C virus NS3 helicase domain. *J. Biol. Chem.* **271**, 24449-24459.
- 21.** Tamura, J., Warrener, P. & Collett, M. (1993) RNA-stimulated NTPase activity associated with the p80 protein of the pestivirus bovine viral diarrhoea virus. *Virology*, **193**, 1-10.